

RESEARCH ARTICLE

(Open Access)

Assessment of the green space of Tirana city and its impact on some public health indicators

ZYDI TEQJA¹; ENKELEJD LEKAJ¹; ZAMIR LIBOHOVA²; NIKOLLAQ BARDHI³¹ Department of Horticulture and Landscape Architecture, Agricultural University of Tirana, Tirana, Albania² USDA-NRCS-National Soil Survey Center, Lincoln, SHBA³ Department of Agronomic Sciences, Agricultural University of Tirana, Tirana, Albania

Abstract

The study of urban green space has become an important topic of research in the last decade. In this study the green area coverage of 11 administrative units of Tirana is evaluated based on the 2015 orthophoto. The total green area and the distribution for administrative units as well as relationships between the amount of green space, suicide, and noise and air pollution over different buffer distances from the monitoring points are evaluated. Over 27% of the area of the city of Tirana is covered by greenery but the distribution is heterogeneous among administrative units. The noise tended to decrease with increasing amount of greenery; however, the trends were not significant. On the other hand, the amount of air pollution decreased significantly with increasing amount of greenery and the strength of the relationship increased with increasing distance from the monitoring points. The relationship between greenery and suicidal rate overall was not significant, except for the most urbanized administrative units; however this based on a limited sample size. The data from this analysis shows trends between the amount of greenery and various health and environmental indicators that could be useful for better planning and management of green spaces in the future, however, more data is needed.

Keywords: Greenery, orthophoto, noise, air pollution, suicide

1. Introduction

The accurate estimation of green area size and distribution is important as it affects urban life and improves health conditions of urban population (6, 13). The definition of the term "green space" can be confusing and biased depending on the measurement or estimation method. This is especially critical for urban areas where the determination of the amount and distribution of green space is challenging due to its heterogeneity. Urban green space is often treated in a general way, but in order to reach accurate and comparable conclusions, a clear definition of green space must be given in each case and what is included in the study must be described. To increase the representation of the important qualities of green spaces, elaborate methods should be used to measure these qualities and green space data with the highest possible resolution are needed to demonstrate the characteristics of urban green space experienced by

people. Such efforts can be seen in some studies that focus on measuring urban green space such as images of green space captured through the Google Street View or using a camera to measure the amount of urban green space visible from perspective of the user (25, 33, and 45). Liang et al, (34) also used the drone to measure the amount of 3D green space.

The choice of measurement method depends on the intent of the researcher and the possibilities for obtaining the data. We can distinguish three groups of methods:

- (i) The first group is based on self-reporting (with surveys given to participants to measure the perceived availability of green space);
- (ii) The second group is based on inventories of areas planted with different vegetation;
- (iii) The third group is based on aerial imagery and geographical maps.

*Corresponding author: Zydi Teqja; E-mail: zteqja@ubt.edu.al

Special Issue of the Conference: Agriculture a Life Science with Roots in Applied Biology; 3-4 Dec. 2020. (Accepted for publication 4.01.2020)

ISSN: 2218-2020, © Agricultural University of Tirana

There are advantages and limitations in determining the most accurate coverage of green space by each group of methods (43). The measurement of green space through questionnaires is based on the study of the population living in the study area. Questionnaires provided to the population in urban areas can be useful when assessing information about green space such as: proximity to home, accessibility, perceptions of quality and current use (28; 31). However, this method requires intensive resources and may not be possible in all studies. Also, the accuracy of the method is affected by the self-reporting of individuals that can result in a biased classification (19). Self-reporting of neighborhood characteristics could be affected by the health conditions of participating individuals. This may be especially true when the outcome includes psychological disorders such as depression (35, 37, and 42).

Measurement of green spaces based on inventories of land planted with vegetation often does not consider the trees and other greenery on the side of roads or urban squares thus removing a significant amount of urban greenery.

Measuring green space using mapping tools is based on Geographic Information Systems (GIS) technology. GIS is a powerful computer mapping technology and analysis that allows large amounts of information to be viewed and analyzed within a geographical context (53). Basically, the GIS program electronically defines large geographical areas to see the physical characteristics of a given land, using satellite imagery (Landsat, CORINE, etc. 30, 32, 50, and 51). GIS along with spatial analytical techniques are widely used to study public health issues (11, 41). Through various methods, GIS can be used to determine the area of land that is classified as green space. These methods include various geographic maps and existing data in the GIS program.

One index that is used to measure the amount of green space in urban areas is the Normalized Difference Vegetation Index (NDVI). NDVI provides objective measurements of healthy vegetation levels using Landsat images that can be applied to almost all land surfaces worldwide (15, 21, and 46). It is widely used in forest and agriculture research. In some researches, NDVI has been used in studies of the links between neighborhood greenery and health (4, 5, 13, 29, 52).

However, the use of NDVI for the definition of green space in urban environments may have some limitations. For example, spatial distribution of

vegetation may be less concentrated and more difficult to detect at high levels of population density, compared to neighborhoods with lower densities (46). Also, an NDVI pixel in an urban setting rarely covers a single homogeneous region (as in the case of the agricultural environment) which makes it difficult to determine the vegetation within that pixel. Another limitation relates to the fact that there are many factors that affect the relationship between NDVI and geographical and physical attributes. These may include atmospheric conditions, image scale, vegetation moisture, soil moisture, general vegetation cover, and changes in soil type (1).

Finally, a major limitation to using NDVI to define green (or "open") spaces is its reliance on color reflection as a method of detection. This can greatly affect the determination of the amount of these areas considering the seasons. For example, during the seasons the leaves normally change color from green to red or yellow, which would result to lower NDVI score (22). This limitation can be addressed to a certain degree by applying the NDVI during late spring and early summer when the green coverage is likely at its peak.

An alternative method to NDVI is the classification of high resolution orthophotos, when available. ArcGIS offers flexibilities for more detailed spatial analysis of green space and pattern distribution.

For example, the high resolution orthoimage can be used to determine the area covered by the green space within a user specified buffer zone around any objects (streets, building, houses, etc.). However, this method depends on the definition of green space and on the manual characterization of land as "green space" and "other space". Based on the definition of green space used the method can be very subjective. Also, the method may require field validation in order to ensure accuracy.

In addition to amount and distribution an important indicator for green space is density. This would require an assessment of the strength of reflection in the case of satellite image or color in case of orthoimages. These are more challenging to quantify. Albania was a rural country until beginning of the new century. The rapid changes that happened during last decade of twentieth century (49) were crowned in 2007 when for the first time the majority of Albanian population lives in urban areas. As developing countries like Albania move toward urban development the disconnectedness from nature is

increasing and it brings in scene the scientific interest in the potential health benefits of nature contact. Though there is evidence to suggest that contact with nature gives a range of health benefits and research recently has yielded substantial evidence on the role of urban green spaces, still there are large gaps in our understanding on this relationship (16, 20).

According to a WHO report (55), mental disorders contribute to many suicides around the world. Over 800 000 people die due to suicide every year and it is the second leading cause of death in 15-29-year-olds (WHO 2014). This is why the research on the benefits of mental health through reduced depression (3, 5, 10, 17, 27, 36, 40, 48, 54); Reduced anxiety (2, 5, 9, 41, 47) and Stress recovery and Suicide prevention (2, 8, 18, 24, 44, 55) seem to be more challenging.

The objective of this study was to: (i) determine the amount and distribution of greenery in the capital city of Tirana (Albania), and (ii) the relationships between the amount of greenery and health and environmental indicators.

2. Materials and Methods

2.1 Data sources

The geo-rectified orthophoto of 2015 for the city of Tirana provided by ASIG (The State Authority for Geospatial Information) was used for this study. The Projection of the ortho image was European Terrestrial Reference System 1989; GRS 1980 Ellipsoid, Lambert Equal Area Projection (ETRS_1989_LAEA) with an 8 cm pixel resolution. The data on the level of noise and pollution for some monitoring points of the city of Tirana were provided through the National Environment Agency and data on the number and causes of death according to the administrative units of the city of Tirana were provided by the General Directorate of Civil Status.

2.2 Data Processing

The ortho image in RGB color space was classified in 5 classes using Iso Cluster Unsupervised Classification in ArcMap 10.5. The 11 polygons administrative units of Tirana were converted to raster using the same resolution and extend (snap) as that of classified image. Administrative units were used as mask to extract the classified image. The area of each

class was calculated in m² and km² then zonal statistics was run to determine the percent green area for each administrative unit. After that, the distribution of green space for 11 units based on squares with 200m sides was estimated by calculating the standard deviation and the coefficient of variation. To assess the impact of greenery on noise and air pollution (particulate matter-PM₁₀ in µg /m³), the green space was calculated for buffer zones of 25m, 50m, 100m, 200m radius from the monitored points for noise level and PM₁₀ in the municipality of Tirana. To better quantify the spatial distribution of green spaces within each administrative unit, the study area was divided into squares of 200x200m and the number of tiles with green spaces was calculated. The coefficient of variation of the distribution of greenery within each unit was determined. These squares were also used to test the accuracy of the green area measurement by means of orthophoto conversion. For this purpose, manual digital measurement was performed in 11 squares with a surface of 200x200m randomly selected (one for each administrative unit).

2.3 Data Analysis

Simple linear regression analysis was conducted to assess the relationships between the amount of greenery and health and environmental indicators. The analysis was performed in Excel for Microsoft Office 365.

3. Results and Discussions

3.1 Amount and distribution characteristics of green space

Figure 1 shows the general view of the orthophoto of the city of Tirana provided by ASIG, while Figure 2 shows the same image after the color classification with the administrative divisions of the urban area of Tirana. The third category of color values in Figure 2 represents the green space. After extracting the classified image it was possible to calculate the area of each class and then zonal statistics was run to determine the percent of green area for each administrative unit. The total area, green spaces and some spatial distribution characteristics for the 11 administrative units of the municipality of Tirana are given in Table 1.



Figure 1. Orthophoto of Tirana 2015 (Source ASIG Albania: <https://geoportal.asig.gov.al/>).




Figure 2. The classification of the ortho image for Tirana city; the borders of Administrative units and the locations of noise and PM10 pollution observations. The third category of color values represents the green space.

Table 1 Spatial distribution characteristics of Tirana green space, life span and suicidal cases by administrative units.

| Administrative Unit | Total Area (km2) | Green Area (km2) | % of Green cover | Standard deviation | Variation coefficient | Life Span (years) | Average Suicidal cases | Suicidal Rate (/1000 deaths) |
|---------------------|------------------|------------------|------------------|--------------------|-----------------------|-------------------|------------------------|------------------------------|
| 1 | 3.06 | 1.22 | 39.9 | 10307 | 65% | 72.9 | 2.7 | 10.3 |
| 2 | 9.58 | 3.98 | 41.5 | 9174 | 56% | 73.9 | 4.2 | 10.1 |
| 3 | 2.01 | 0.4 | 19.9 | 6000 | 78% | 73.0 | 2.0 | 8.2 |
| 4 | 4.92 | 1.29 | 26.2 | 5850 | 56% | 71.8 | 0.8 | 2.8 |
| 5 | 2.87 | 0.46 | 16.0 | 3940 | 65% | 74.9 | 2.2 | 5.0 |
| 6 | 5.48 | 1.67 | 30.5 | 7894 | 71% | 71.6 | 3.2 | 9.6 |
| 7 | 3.06 | 0.48 | 15.7 | 4176 | 70% | 75.2 | 2.0 | 5.7 |

Tirana Green space and its impact on public health

| | | | | | | | | |
|----|------|------|------|------|-----|------|-----|-----|
| 8 | 1.95 | 0.28 | 14.4 | 3311 | 58% | 74.8 | 1.7 | 7.1 |
| 9 | 2.82 | 0.55 | 19.5 | 6094 | 72% | 73.8 | 1.5 | 4.8 |
| 10 | 0.77 | 0.11 | 14.3 | 2567 | 48% | 76.7 | 1.3 | 6.5 |
| 11 | 5.21 | 1.12 | 21.5 | 5253 | 60% | 71.2 | 2.3 | 8.5 |

Figure 3 shows what percentage of the green area for each administrative unit. A significant part of the green area consists of greenery from trees along the streets, which is an important part of urban greenery. About 27.7% of the territory of the city of Tirana is covered by greenery with a somewhat heterogeneous distribution among the administrative units. Thus, the amount of green space varies from about 14.4% (administrative unit no. 8) to about 40% (administrative units 1 and 2). The administrative units 1 and 2 have only recently expanded to include former rural areas where the density of construction is still low.

To validate our model and to better understand, what the distribution of green spaces is in each administrative unit, the city territory was divided into squares with 200m sides. These squares were used to test the accuracy of the green area measurement by means of orthophoto conversion. For this purpose, manual digital measurement was performed in 11 squares with a surface of 200x200m randomly selected (one for each administrative unit). Figure 4 shows the relationship between the values measured through the orthophoto conversion model and the values measured manually. The strong correlation

between these values is clear, which indicates the high accuracy of the measurement by our model. Note that all manual measurements are slightly lower than those by model. This may be due to the fact that by manual measurement it is difficult to include every green pixel in the measurement. This is evident in the fact that the measurements are almost equal when the green area is large and compact. It was therefore judged that there is no need to make correction to the model measurements.

This was followed by the analysis of the distribution of greenery within each unit by means of the coefficient of variation. These spatial distribution characteristics of the green space within each administrative unit are given in Table 1. The analysis shows that administrative units 3, 6 and 9 have the highest coefficient of variation, which means that they have the most heterogeneous distribution of green space. Units 2, 4 and 10, on the other hand, have a lower coefficient of variation, so they have a more uniform distribution of green space. Likely, the residents of these units have more equal opportunities to benefit from greenery.




Figure 3. Green space in percentage of the total area by administrative units.




Figure 4. The relationship between manual and model measurements of green cover

3.2 Impact of green space on noise

To see the impact of greenery on noise levels we analyzed 15 monitored points for noise level (Figure 1) and calculated the green area in buffer zones with

radiuses 25m, 50m, 100m, and 200m. The data are shown in Table 2.

Table 2. Green space amount in m2 for varying buffer distances from 15 Tirana city monitoring points. (Source of noise data: Reports of the National Environmental Agency of 2016).

| No | Monitoring points | Green cover for different buffer zones | | | | Year 2015 | |
|----|----------------------|--|------|-------|-------|-----------|-------|
| | | 25m | 50m | 100m | 200m | Day | Night |
| 1 | American Bank | 25 | 258 | 2010 | 13202 | 63.47 | 62.83 |
| 2 | Farmacia 10 | 85 | 664 | 3148 | 16995 | 67.63 | 53.44 |
| 3 | 21 Dhjetori | 299 | 685 | 2663 | 11163 | 68.41 | 61.05 |
| 4 | Vasil Shanto | 422 | 1888 | 6672 | 19929 | 66.39 | 54.89 |
| 5 | Police Directorate | 501 | 1360 | 4239 | 14546 | 64.76 | 53.67 |
| 6 | Selvia | 39 | 221 | 2028 | 11199 | 65.39 | 58.72 |
| 7 | Laprake | 92 | 221 | 777 | 7698 | 67.69 | 60.03 |
| 8 | Palace of Congresses | 159 | 879 | 6721 | 33944 | 64.27 | 54.47 |
| 9 | Unknown Partizan | 258 | 777 | 3538 | 19096 | 67.35 | 57.34 |
| 10 | Shkolla Bashkuar | 117 | 1105 | 5591 | 19523 | 66.21 | 53.47 |
| 11 | Dinamo Stadion | 238 | 1088 | 4319 | 16273 | 59.45 | 58.54 |
| 12 | Elbasani Street | 226 | 1011 | 4350 | 17927 | 71.16 | 61.43 |
| 13 | Skenderbeg | 41 | 1031 | 10254 | 26707 | 74.25 | 62.49 |
| 14 | Train Station | 96 | 421 | 3331 | 13567 | 64.04 | 61.63 |
| 15 | Edith Durham school | 316 | 1500 | 3938 | 15644 | 61.69 | 58.07 |

The graphs in Figures 4 and 5 show the relationship between the green area and the noise level at night for 15 points at 25 and 50 m distance. These figures show a tendency to reduce noise with increasing greenery,

but it should be noted that the relationship is weak and statistically unverified, so this issue needs more studies and perhaps better data quality.




Figure 4. Relationship between overnight noise level and amount of greenery within a 25m radius of the monitoring point.




Figure 5 Relationship between overnight noise level and amount of greenery within a 50m radius of the monitoring point

3.3 Impact of green space on air pollution (PM10)

It was possible to find data on the level of pollution by PM10 particles for 11 monitoring points (Figure 1).

Monitoring points for PM10 $\mu\text{g} / \text{m}^3$ level and green area in buffer zones with radiuses 50m, 100m, 200m are given in table 3.

Table 3 Monitoring points for PM10 and green area for different buffer zones.

| No | Monitoring points | Green cover for different buffer zones | | | PM10 $\mu\text{g}/\text{m}^3$ |
|----|--------------------|--|------|-------|-------------------------------|
| | | 50m | 100m | 200m | |
| 1 | Farmacia 10 | 631 | 2803 | 16576 | 40.62 |
| 2 | Iranian Clinic | 800 | 3197 | 17980 | 34.40 |
| 3 | Paris Commune | 90 | 720 | 8165 | 28.59 |
| 4 | 21 Dhjetori | 196 | 2478 | 10743 | 73.65 |
| 5 | Koco Glozheni M | 1779 | 5501 | 16696 | 31.20 |
| 6 | Palace with arrows | 486 | 3871 | 16421 | 48.68 |
| 7 | Ali Demi St. | 1367 | 4723 | 16342 | 45.60 |
| 8 | Elbasani Street | 1602 | 5128 | 18337 | 37.55 |
| 9 | Dynamo Stadium | 902 | 3972 | 18242 | 36.72 |
| 10 | Train Station | 567 | 2374 | 13829 | 51.25 |
| 11 | Zogu Zi | 604 | 1506 | 6734 | 71.18 |

Source PM10 data: ISHP

Figure 6 shows the association of green spaces for different buffer zones from the monitoring point with the level of PM10 particles. It can be seen that the strength of the green area impact on PM10 increases

with distance from the monitoring point represented by the buffer size. Increasing the green area leads to significant reductions in pollution.




Figure 6. Impact of green space on PM10 for (A) 50m Buffer; (B) 100m Buffer; and (C) 200m Buffer.

Figures 6 a, b and c show that the strength of the green area impact on PM10 increases with distance from the monitoring point represented by the buffer size. Increasing the green area leads to significant reductions in pollution. The impact of greenery on air pollution has been discussed in different studies. Some of them consider the management of urban tree canopy cover as a viable strategy to improve air quality and help meet clean air standards (38, 39).

3.4 Greenery and suicides

The impact of greenery on mental health problems has been highlighted by several authors (references). In this study we analyzed the relationship between the number of suicides (as an extreme case of mental health) and the amount of greenery by administrative units (Figure 7).




Figure 7 (A) Greenery and summer suicides for Tirana city; (B) Greenery and suicides for the May-October period for the most urbanized area of Tirana city.

If we perform a simple analysis for this relationship it seems that there is no relationship between these parameters. But if we focus only on the warm period of the year (in fact this is the period when the need for greenery is most felt) we see the emergence of a trend in this relationship (Figure 7 A). If we separate from this relationship the administrative units which in their territory have large rural areas and focus on the real urban area, so if we exclude the administrative units 1, 2, 6 and 11 then a clear connection is found between the green area and the number of suicides. So there seems to be a high and significant relationship between the amount of green and suicidal rates occurring during May-October period for the most urbanized administrative units (Figure 7B).

This situation could be explained as follows:

1. The administrative units 1, 2, 6 and 11 in recent years have included in their territories many areas that were previously outside the city and continue to have a pronounced rural character and it does not make sense to analyze the impact of greenery in rural areas where greenery is abundant.
2. In these areas there is a great change of population due to internal migration. So a large part of their population has not been there all their life but have come the last two decades.
3. As a consequence of the above two facts, these units also have important socio-economic differences with the most urbanized part of the city, differences

that make the studied connection much more complex and justify their separation.

According to different studies flora and fauna in urban green space can reduce blood pressure, depression and boost immune function. Vegetation in urban green space can also reduce health risks by mitigating air pollution and reducing the effect of hot islands in urban areas (28) as well as regulating other aspects of the microclimate.

Also natural views and sounds created by flora and fauna can increase parasympathetic activation, which can reduce metabolism and promote relaxed and calm feelings (23). Other structures in urban green space such as open space and playgrounds can encourage health behaviors through physical activity and social interactions, which have resulted in positive health indicators (12, 14).

It is very likely that the impact on depression and suicide is both from the effect that green space have on pollution reduction and from the direct effect of greenery. According to a recent study (7) exposure to air pollution is associated with an increased risk of depression and suicide. This analysis reviewed 25 studies published by the end of 2017 and found that someone living for at least six months in an area with twice the limit recommended by World Health Organization for fine particles, PM_{2.5}, there would be approximately a 10 percent risk of increased depression. For suicides a link was found between short-term exposures to PM₁₀ contamination. Thus exposure to 10 µg / m³ of PM₁₀ over a three-day period was associated with a 2 percent higher risk of suicide (7).

Although the data shows a relationship between suicidal rate and the amount of green space especially for the May-October months, the results from this analysis need to be viewed critically. First, there is a very limited number of data. Second, there may be other underlying reasons for the observed suicidal rates and their relationship with the amount of green space. So, more studies and data are needed for reaching a definite conclusion.

4. Conclusions

Orthophotos provide very high resolution images making them very useful to measure and study green cover in urban areas where satellite resources like CORINE Land Cover cannot identify green spaces because of low resolution. In this study the orthophoto with 8cm resolution is used to measure and study the green cover of Tirana city. More than 27% of the area

of the city of Tirana is covered by green vegetation. There is a heterogeneous distribution of green space between administrative units varying from 15% to 40%.

The relationships between environmental indicators (noise and air pollution) and the amount of green cover, though not all significant highlight in general the importance of green space for a healthy life. The relationship between suicidal rate and amount of green space, though intriguing should be interpreted with caution and merits further research.

The results from this study could potentially aide in better planning and management of green spaces in the future.

5. References

1. Abuzar M, Sheffield K, Whitfield D, O'Connell M, McAllister A: **Comparing inter sensor NDVI for the analysis of horticulture crops in south-eastern Australia.** American Journal of Remote Sensing. 2014;2(1):1-9.
2. Annerstedt M, Ostergren PO, Björk J, Grahn P, Skärbäck E, Währborg P: **Green qualities in the neighbourhood and mental health - results from a longitudinal cohort study in Southern Sweden.** BMC Public Health. 2012 May 8;12:337. doi: 10.1186/1471-2458-12-337. PMID: 22568888; PMCID: PMC3443019.
3. Astell-Burt T, Feng X: **Association of Urban Green Space With Mental Health and General Health Among Adults in Australia.** JAMA Netw Open. 2019;2(7):e198209. doi:10.1001/jamanetworkopen.2019.8209
4. Bell JF, Wilson J S, Liu G C: **Neighborhood greenness and 2-year changes in body mass index of children and youth.** Am J Prev Med. 2008;35(6):547-53
5. Beyer KM, Kaltenbach A, Szabo A, Bogar S, Nieto FJ, Malecki KM: **Exposure to neighborhood green space and mental health: evidence from the survey of the health of Wisconsin.** International journal of environmental research and public health. 2014;11(3):3453-72.
6. Beyer KMM, Kaltenbach A, Szabo A, Bogar S, Nieto F J, Malecki KM: **Exposure to neighborhood green space and mental health: Evidence from the survey of the health of Wisconsin.** Int J Environ Res

- Public Health 2014;11(3): 3453– 3472, <https://doi.org/10.3390/ijerph110303453>.
7. Braithwaite I, Zhang S, Kirkbride JB, Osborn DPJ, Hayes JF: **Air pollution (particulate matter) exposure and associations with depression, anxiety, bipolar, psychosis and suicide risk: a systematic review and meta-analysis.** Environ Health Perspect. 2019;127(12):126002.
8. Bratman G N, Hamilton J P, Hahn K S, Daily G C, Gross J J: **Nature experience reduces rumination and subgenual prefrontal cortex activation.** Proc Natl Acad Sci USA 2015, 112(28):8567–8572.
9. Bratman GN, Hamilton JP, and Daily G C: **The impacts of nature experience on human cognitive function and mental health.** Ann NY Acad Sci 2012, 1249:118–136, PMID: 22320203, <https://doi.org/10.1111/j.1749-6632.2011.06400.x>.
10. Cohen-Cline H, Turkheimer E, Duncan GE: **Access to green space, physical activity and mental health: a twin study.** J Epidemiol Community Health 2015, 69(6): 523–529, PMID: 25631858, <https://doi.org/10.1136/jech-2014-204667>.
11. Cromley E, McLafferty S: **Public participation GIS and community health.** GIS and public health. 2012:411-22.
12. Dadvand P, Bartoll X, Basagaña X, Dalmau-Bueno A, Martinez D, Ambros A, Cirach M, Triguero-Mas M, Gascon M, Borrell C, Nieuwenhuijsen M J, Green spaces and General Health: **Roles of mental health status, social support, and physical activity,** Environment International, Volume 91, 2016, Pages 161-167, ISSN 0160-4120, <https://doi.org/10.1016/j.envint.2016.02.029>.
13. Dadvand P, Sunyer J, Basagaña X, Ballester F, Lertxundi A, Fernández-Somoano A, et al. **Surrounding greenness and pregnancy outcomes in four Spanish birth cohorts.** Environ Health Perspect. 2012;120(10):1481-7
14. de Vries S, van Dillen SM, Groenewegen PP, Spreeuwenberg P. **Streetscape greenery and health: stress, social cohesion and physical activity as mediators.** Soc Sci Med. 2013 Oct;94:26-33. doi: 10.1016/j.socscimed.2013.06.030. Epub 2013 Jul 3. PMID: 23931942.
15. Eastman JR, Sangermano F, Machado EA, Rogan J, Anyamba A. **Global trends in seasonality of normalized difference vegetation index (NDVI), 1982–2011.** Remote Sensing. 2013;5(10):4799-818.
16. Frumkin, H., Bratman, N., Breslow S.J., Cochran, B., Kahn, P.H., Lawler, J.J., Levin, P.S., Tandon, P.S., Varanasi, U., Wolf, K.L. and Wood, S.A. (2017) **Nature contact human health; A research agenda.** Environ. Health Perspect 125: Published:31 July 2017 CID: 075001 <https://doi.org/10.1289/EHP1663>
17. Gascon M, Triguero-Mas M, Martinez D, Dadvand P, Fornes J, Plasencia A, et al.: **Mental health benefits of long-term exposure to residential green and blue spaces: a systematic review.** Int J Environ Res Public Health 2015, 12(4): 4354– 4379, PMID: 25913182, <https://doi.org/10.3390/ijerph120404354>.
18. Helbich, M., de Beurs, D., Kwan, M. P., O'Connor, R. C., & Groenewegen, P. P: **Natural environments and suicide mortality in the Netherlands: a cross-sectional, ecological study.** The Lancet. Planetary health, 2018, 2(3), e134–e139. [https://doi.org/10.1016/S2542-5196\(18\)30033-0](https://doi.org/10.1016/S2542-5196(18)30033-0)
19. Hillsdon M, Panter J, Foster C, Jones A: **The relationship between access and quality of urban green space with population physical activity.** Public health. 2006;120(12):1127-32.
20. Hossain, M., Sultana, A., Ma, P., Fan, Q., Sharma, R., Purohit, N., & Sharmin, D. F: **Effects of natural environment on mental health: an umbrella review of systematic reviews and meta-analyses.** PsyArXiv, 8 Jan. 2020; <https://doi.org/10.31234/osf.io/4r3mh>
21. Hysa A, Teqja Z: **Counting fuel properties as input in the wildfire spreading capacities of vegetated surfaces: case of Albania.** Notulae Botanicae Horti Agrobotanici Cluj-Napoca 2020, 48(3):1667-1682; DOI:10.15835/nbha48311994.

22. Jackson RD, Huete AR: **Interpreting vegetation indices**. *Prev Vet Med*. 1991;11(3-4):185-200.
23. James AC, James G, Cowdrey FA, Soler A, Choke A: **Cognitive behavioural therapy for anxiety disorders in children and adolescents**. *Cochrane Database of Systematic Reviews* 2015, Issue 2. Art. No.: CD004690. DOI: 10.1002/14651858.CD004690.pub4.
24. Jiang, B., Li, D., Larsen, L., & Sullivan, W. C: **A Dose-Response Curve Describing the Relationship Between Urban Tree Cover Density and Self-Reported Stress Recovery**. *Environment and Behavior*, 2014, 48(4), 607-629.
<https://doi.org/10.1177/0013916514552321>
25. Kang J, Körner M, Wang Y, Taubenböck H, Zhu X: **Building instance classification using street view images**, *ISPRS Journal of Photogrammetry and Remote Sensing*, Volume 145, Part A, 2018, Pages 44-59, ISSN 0924-2716, <https://doi.org/10.1016/j.isprsjprs.2018.02.006>.
26. Kardan, O, Gozdyra, P, Misic B. et al.: **Neighborhood greenspace and health in a large urban center**. *Sci Rep* 5, 11610 (2015).
<https://doi.org/10.1038/srep11610>
27. Korpela Kalevi, M., Stengård Eija and Jussila Pia: **Nature Walks as a Part of Therapeutic Intervention for Depression**. *Ecopsychology* 8 (2016): 8-15.
28. Kuo FE, Sullivan WC: **Environment and crime in the inner city does vegetation reduce crime?** *Environment and behavior*. 2001;33(3):343-67.
29. Lachowycz K, Jones A: **Greenspace and obesity: a systematic review of the evidence**. *Obesity reviews*. 2011;12(5):e183-e9.
30. Lekaj E, Teqja Z: **Investigation of green space changes in Tirana-Durres region**. Third International Conference "Challenges in Biotechnological and Environmental Approaches", Agricultural University of Tirana, ALBANIA, April 23 - 25, 2019. *Albanian J. Agric. Sci.* (Special Edition)
31. Lekaj E, Teqja Z: **Users evaluation for public parks: influences of location, season, gender and age**. *European Academic Research*, 2019, VII(1), 603-624.
32. Lekaj E, Teqja Z, Bani A: **The dynamics of land cover changes and the impact of climate change on ultramafic areas of Albania**. *Periodico di Mineralogia* 2019, 88, 223-234. (IF 1.417)
33. Li X, Chuanrong Zhang, Weidong Li, Robert Ricard, Qingyan Meng, Weixing Zhang: **Assessing street-level urban greenery using Google Street View and a modified green view index**, *Urban Forestry & Urban Greening*, Volume 14, Issue 3, 2015, Pages 675-685, ISSN 1618-8667, <https://doi.org/10.1016/j.ufug.2015.06.006>.
34. Liang H, Weizheng Li, Qingping Zhang, Wen Zhu, Di Chen, Jie Liu, Ting Shu: **Using unmanned aerial vehicle data to assess the three-dimension green quantity of urban green space: A case study in Shanghai, China**, *Landscape and Urban Planning*, Volume 164, 2017, Pages 81-90, ISSN 0169-2046, <https://doi.org/10.1016/j.landurbplan.2017.04.006>.
35. Mair CF, Roux AVD, Galea S: **Are neighborhood characteristics associated with depressive symptoms? A critical review**. *J Epidemiol Community Health*. 2008;jech. 2007.066605.
36. McEachan R R, Prady S L, Smith G, Fairley L, Cabieses B, Gidlow C, et al.: **The association between green space and depressive symptoms in pregnant women: Moderating roles of socio economic status and physical activity**. *J Epidemiol Community Health* 2016, 70(3): 253–259, PMID: 26560759, <http://dx.doi.org/10.1136/jech-2015-205954>
37. Mitchell R, Astell-Burt T, Richardson EA: **A comparison of green space indicators for epidemiological research**. *J Epidemiol Community Health*. 2011 February 4, 2011.
38. Nowak, D. J., Crane, D. E. & Stevens, J. C: **Air pollution removal by urban trees and shrubs in the United States**. *Urban forestry & urban greening* 4, 115–123 (2006).
39. Nowak, D. J., Hirabayashi, S., Bodine, A. & Greenfield, E: **Tree and forest effects on air quality and human health in the United**

- States.** Environmental Pollution 193, 119–129 (2014).
40. Nutsford D, Pearson A L, Kingham S: **An ecological study investigating the association between access to urban green space and mental health.** Public Health 2013, 127(11):1005–1011, PMID: 24262442, <https://doi.org/10.1016/j.puhe.2013.08.016>
41. Nykiforuk CI, Flaman LM: **Geographic information systems (GIS) for health promotion and public health: a review.** Health promotion practice. 2011;12(1):63-73.
42. O'campo P, Kub J, Woods A, Garza M, Jones AS, Gielen AC, et al.: **Depression, PTSD, and comorbidity related to intimate partner violence in civilian and military women.** Brief treatment and crisis Intervention. 2006; 6(2):99.
43. Patel, D., Block, R., Chapman, B., Korfmacher, K., & Wijngaarden, E.V: **Green space and mental health symptoms in a cardiac rehabilitation population.** Indoor and Built Environment, 2019, 28, 1431 - 1440.
44. Pollock N. J: **Place, the Built Environment, and Means Restriction in Suicide Prevention.** International journal of environmental research and public health, 2019, 16(22), 4389. <https://doi.org/10.3390/ijerph16224389>
45. Quynh C. Nguyen, Sahil Khanna, Pallavi Dwivedi, Dina Huang, Yuru Huang, Tolga Tasdizen, Kimberly D. Brunisholz, Feifei Li, Wyatt Gorman, Thu T. Nguyen, Chengsheng Jiang: **Using Google Street View to examine associations between built environment characteristics and U.S. health outcomes,** Preventive Medicine Reports, Volume 14, 2019, 100859, ISSN 2211-3355
46. Rhew IC, Vander Stoep A, Kearney A, Smith NL, Dunbar MD: **Validation of the normalized difference vegetation index as a measure of neighborhood greenness.** Ann Epidemiol. 2011;21(12):946-52.
47. Song C, Ikei H, Igarashi M, Takagaki M, Miyazaki Y: **Physiological and psychological effects of a walk in urban parks in fall.** Int J Environ Res Public Health 12(11):14216–14228, PMID: 26569271, <https://doi.org/10.3390/ijerph121114216>.
48. Taylor, Mark S., Wheeler, Benedict W., White, Mathew P, Economou, Theodoros, and Osborne, Nicholas J: **Research note: urban street tree density and antidepressant prescription rates - a cross-sectional study in London, UK.** Landscape and Urban Planning 2015, 136 174-179. <https://doi.org/10.1016/j.landurbplan.2014.12.005>, <https://doi.org/10.1016/j.landurbplan.2014.12.005>.
49. Teqja Z, Beka I, Shkreli E: **Albanian agriculture- dramatic changes from a very centralized economy to free market.** A strategy for future development. Published in “Medit” no. 1, 2000, pages 21-29.
50. Teqja Z, Lekaj E.: **Land cover changes evaluation of Tirana region.** Albanian j. agric. sci. ISSN: 2218-2020, Volume 16, Special issue (2017).
51. Teqja Z, Lekaj E, Bardhi N: **Using land cover to calculate some nature indicators of the Republic of Albania.** A.J.A.S Vol 16 (1) pp. 19-27. Publ. Date: 31 March 2017.
52. Villeneuve PJ, Jerrett M, Su JG, Burnett RT, Chen H, Wheeler AJ, et al.: **A cohort study relating urban green space with mortality in Ontario, Canada.** Environ Res. 2012;115:51-8.
53. Vine MF, Degnan D, Hanchette C: **Geographic information systems: their use in environmental epidemiologic research.** Environ Health Perspect. 1997;105(6):598. 200
54. White MP, Alcock I, Wheeler B W, Depledge M H: **Would you be happier living in greener urban area? A fixed-effects analysis of panel data.** Psychol Sci 2013, 24(6):920–928, <https://doi.org/10.1177/0956797612464659>.
55. World Health Organization. **Preventing Suicide: A Global Imperative;** 9241564776; WHO: Geneva, Switzerland, 2014; Available online: http://www.who.int/mental_health/suicide-prevention/world_report_2014/en/